

Aricept Evess tablets are indicated for the treatment of MILD, MODERATE and SEVERE dementia in Alzheimer's Disease.

■ EARLIER

Earlier intervention preserves more function¹⁻³

■ HIGHER

Higher the dose, better the response⁴⁻⁶

■ LONGER

Longer the duration of treatment, better the long-term outcome⁷⁻⁹

■ BETTER

Better efficacy across all of the Alzheimer's Disease domains such as cognition, global function, neuropsychiatry symptoms and quality of life¹⁰⁻¹²

References

1. L. S. Galvin et al. *Alzheimer's & Dementia* 2014; 10: 3. 2. Galvin L et al. *Alzheimer's & Dementia* 2014; 10: 3. 3. Galvin L et al. *Alzheimer's & Dementia* 2014; 10: 3. 4. Galvin L et al. *Alzheimer's & Dementia* 2014; 10: 3. 5. Galvin L et al. *Alzheimer's & Dementia* 2014; 10: 3. 6. Galvin L et al. *Alzheimer's & Dementia* 2014; 10: 3. 7. Galvin L et al. *Alzheimer's & Dementia* 2014; 10: 3. 8. Galvin L et al. *Alzheimer's & Dementia* 2014; 10: 3. 9. Galvin L et al. *Alzheimer's & Dementia* 2014; 10: 3.

Additional Information on Aricept Evess

Aricept Evess (donepezil HCl) tablets are indicated for the treatment of mild, moderate and severe dementia in Alzheimer's Disease. They are contraindicated in patients with known hypersensitivity to donepezil or any of the components of the formulation. Aricept Evess should be used with caution in patients with a history of seizures, as donepezil may lower the seizure threshold. Aricept Evess should be used with caution in patients with a history of bradycardia, as donepezil may increase the risk of bradycardia. Aricept Evess should be used with caution in patients with a history of hypotension, as donepezil may increase the risk of hypotension. Aricept Evess should be used with caution in patients with a history of gastrointestinal disorders, as donepezil may increase the risk of gastrointestinal disorders. Aricept Evess should be used with caution in patients with a history of urinary disorders, as donepezil may increase the risk of urinary disorders. Aricept Evess should be used with caution in patients with a history of respiratory disorders, as donepezil may increase the risk of respiratory disorders. Aricept Evess should be used with caution in patients with a history of liver disorders, as donepezil may increase the risk of liver disorders. Aricept Evess should be used with caution in patients with a history of kidney disorders, as donepezil may increase the risk of kidney disorders. Aricept Evess should be used with caution in patients with a history of blood disorders, as donepezil may increase the risk of blood disorders. Aricept Evess should be used with caution in patients with a history of eye disorders, as donepezil may increase the risk of eye disorders. Aricept Evess should be used with caution in patients with a history of ear disorders, as donepezil may increase the risk of ear disorders. Aricept Evess should be used with caution in patients with a history of nose disorders, as donepezil may increase the risk of nose disorders. Aricept Evess should be used with caution in patients with a history of throat disorders, as donepezil may increase the risk of throat disorders. Aricept Evess should be used with caution in patients with a history of mouth disorders, as donepezil may increase the risk of mouth disorders. Aricept Evess should be used with caution in patients with a history of skin disorders, as donepezil may increase the risk of skin disorders. Aricept Evess should be used with caution in patients with a history of hair disorders, as donepezil may increase the risk of hair disorders. Aricept Evess should be used with caution in patients with a history of nail disorders, as donepezil may increase the risk of nail disorders. Aricept Evess should be used with caution in patients with a history of teeth disorders, as donepezil may increase the risk of teeth disorders. Aricept Evess should be used with caution in patients with a history of jaw disorders, as donepezil may increase the risk of jaw disorders. Aricept Evess should be used with caution in patients with a history of neck disorders, as donepezil may increase the risk of neck disorders. Aricept Evess should be used with caution in patients with a history of shoulder disorders, as donepezil may increase the risk of shoulder disorders. Aricept Evess should be used with caution in patients with a history of arm disorders, as donepezil may increase the risk of arm disorders. Aricept Evess should be used with caution in patients with a history of hand disorders, as donepezil may increase the risk of hand disorders. Aricept Evess should be used with caution in patients with a history of wrist disorders, as donepezil may increase the risk of wrist disorders. Aricept Evess should be used with caution in patients with a history of forearm disorders, as donepezil may increase the risk of forearm disorders. Aricept Evess should be used with caution in patients with a history of elbow disorders, as donepezil may increase the risk of elbow disorders. Aricept Evess should be used with caution in patients with a history of upper arm disorders, as donepezil may increase the risk of upper arm disorders. Aricept Evess should be used with caution in patients with a history of lower arm disorders, as donepezil may increase the risk of lower arm disorders. Aricept Evess should be used with caution in patients with a history of forearm disorders, as donepezil may increase the risk of forearm disorders. Aricept Evess should be used with caution in patients with a history of hand disorders, as donepezil may increase the risk of hand disorders. Aricept Evess should be used with caution in patients with a history of wrist disorders, as donepezil may increase the risk of wrist disorders. Aricept Evess should be used with caution in patients with a history of forearm disorders, as donepezil may increase the risk of forearm disorders. Aricept Evess should be used with caution in patients with a history of elbow disorders, as donepezil may increase the risk of elbow disorders. Aricept Evess should be used with caution in patients with a history of upper arm disorders, as donepezil may increase the risk of upper arm disorders. Aricept Evess should be used with caution in patients with a history of lower arm disorders, as donepezil may increase the risk of lower arm disorders.